

anti- ASGR2 antibody

Product Information

Catalog No.: FNab00636

Size: 100μg Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

Clone ID: None IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

Background

Mediates the endocytosis of plasma glycoproteins to which the terminal sialic acid residue on their complex carbohydrate moieties has been removed. The receptor recognizes terminal galactose and N-acetylgalactosamine units. After ligand binding to the receptor, the resulting complex is internalized and transported to a sorting organelle, where receptor and ligand are disassociated. The receptor then returns to the cell membrane surface.

Immunogen information

Immunogen: asialoglycoprotein receptor 2

Synonyms: ASGP R, ASGP R 2, ASGPR 2, ASGR2, asialoglycoprotein receptor 2,

CLEC4H2, Hepatic lectin H2, HL 2, Hs.1259, L H2

Observed MW: 39 kDa UniprotID: P07307

Application

Reactivity: Human

Tested Application: ELISA, WB, IHC, IP

Wuhan Fine Biotech Co., Ltd.


B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)

Tel:(0086)027-87384275 Fax: (0086)027-87800889 www.fn-test.com


1



Recommended dilution: WB: 1:200-1:1000; IP: 1:200-1:1000; IHC: 1:50-1:500 Image:




Immunohistochemistry of paraffin-embedded human liver tissue slide using FNab00636(ASGR2 Antibody) at dilution of 1:200



IP Result of anti-ASGR2 (IP:FNab00636, 4ug; Detection:FNab00636 1:300) with L02 cells lysate 1800ug.





human liver tissue were subjected to SDS PAGE followed by western blot with FNab00636(ASGR2 antibody) at dilution of 1:400