

anti- BBS9 antibody

Product Information

Catalog No.: FNab00825

Size: 100μg Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

Clone ID: None IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

Background

The BBSome complex is thought to function as a coat complex required for sorting of specific membrane proteins to the primary cilia. The BBSome complex is required for ciliogenesis but is dispensable for centriolar satellite function. This ciliogenic function is mediated in part by the Rab8 GDP/GTP exchange factor, which localizes to the basal body and contacts the BBSome. Rab8(GTP) enters the primary cilium and promotes extension of the ciliary membrane. Firstly the BBSome associates with the ciliary membrane and binds to RAB3IP/Rabin8, the guanosyl exchange factor(GEF) for Rab8 and then the Rab8-GTP localizes to the cilium and promotes docking and fusion of carrier vesicles to the base of the ciliary membrane. Required for proper BBSome complex assembly and its ciliary localization.

Immunogen information

Immunogen: Bardet-Biedl syndrome 9

Synonyms: PTHB1
Observed MW: 60 kDa
UniprotID: Q3SYG4

Application

1

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)

Tel:(0086)027-87384275 Fax: (0086)027-87800889 www.fn-test.com




Reactivity: Human, Mouse


Tested Application: ELISA, IHC, IF, WB, IP

Recommended dilution: WB: 1:1000-1:6000; IP: 1:500-1:3000; IHC: 1:50-1:500; IF: 1:50-1:500

Image:




Immunohistochemistry of paraffin-embedded human liver using FNab00825(BBS9 antibody) at dilution of 1:50



IP Result of anti-BBS9 (IP:FNab00825, 4ug; Detection:FNab00825 1:500) with mouse testis tissue lysate 4000ug.

Wuhan Fine Biotech Co., Ltd.





mouse testis tissue were subjected to SDS PAGE followed by western blot with FNab00825(BBS9 antibody) at dilution of 1:300