

anti- CD2 antibody

Product Information

Catalog No.:	FNab01438
Size:	100µg
Form:	liquid
Purification:	Immunogen affinity purified
Purity:	≥95% as determined by SDS-PAGE
Host:	Rabbit
Clonality:	polyclonal
Clone ID:	None
IsoType:	IgG
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months (Avoid repeated freeze / thaw cycles.)

Background

CD2 interacts with lymphocyte function-associated antigen(LFA-3) and CD48/BCM1 to mediate adhesion between T-cells and other cell types. CD2 is implicated in the triggering of T-cells, the cytoplasmic domain is implicated in the signaling function.

Immunogen information

Immunogen:	CD2 molecule
Synonyms:	SRBC
Observed MW:	55 kDa
Uniprot ID :	P06729

Application

Reactivity:	Human, Mouse, Rat
Tested Application:	ELISA, IHC, WB
Recommended dilution:	WB: 1:500-1:2000; IHC: 1:20-1:200
Image:	


Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 Gaoxin Ave. East Lake High-Tech Development Zone. Wuhan, Hubei, China(430206)


Tel : (0086)027-87384275

Fax: (0086)027-87800889

www.fn-test.com



Immunohistochemistry of paraffin-embedded human tonsillitis tissue slide using FNab01438(CD2 Antibody) at dilution of 1:200 Heat mediated antigen retrieved with Tris-EDTA buffer, pH9.0



Jurkat cells were subjected to SDS PAGE followed by western blot with FNab01438(CD2 Antibody) at dilution of 1:3000