

anti- CDC26 antibody

Product Information

Catalog No.: FNab01524

Size: 100μg Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

Clone ID: None IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

Background

Component of the anaphase promoting complex/cyclosome(APC/C), a cell cycle-regulated E3 ubiquitin ligase that controls progression through mitosis and the G1 phase of the cell cycle. The APC/C complex acts by mediating ubiquitination and subsequent degradation of target proteins: it mainly mediates the formation of 'Lys-11'-linked polyubiquitin chains and, to a lower extent, the formation of 'Lys-48'-and 'Lys-63'-linked polyubiquitin chains. May recruit the E2 ubiquitin-conjugating enzymes to the complex.

Immunogen information

Immunogen: cell division cycle 26 homolog(S. cerevisiae)

Synonyms: ANAPC12, C9orf17

Observed MW: 10 kDa UniprotID: Q8NHZ8

Application

Reactivity: Human, Mouse, Rat Tested Application: ELISA, WB, IHC

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)


Tel:(0086)027-87384275 Fax: (0086)027-87800889 www.fn-test.com

1




Recommended dilution: WB: 1:500-1:2000; IHC: 1:20-1:200

Image:



Immunohistochemistry of paraffin-embedded mouse brain tissue slide using FNab01524(CDC26 Antibody) at dilution of 1:50



HeLa cells were subjected to SDS PAGE followed by western blot with FNab01524(CDC26 Antibody) at dilution of 1:300