

# anti- COPS4 antibody

#### **Product Information**

Catalog No.: FNab01872

Size:  $100\mu g$  Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

Clone ID: None IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

### **Background**

Component of the COP9 signalosome complex(CSN), a complex involved in various cellular and developmental processes. The CSN complex is an essential regulator of the ubiquitin(Ubl) conjugation pathway by mediating the deneddylation of the cullin subunits of SCF-type E3 ligase complexes, leading to decrease the Ubl ligase activity of SCF-type complexes such as SCF, CSA or DDB2. Also involved in the deneddylation of non-cullin subunits such as STON2. The complex is also involved in phosphorylation of p53/TP53, c-jun/JUN, IkappaBalpha/NFKBIA, ITPK1, IRF8/ICSBP and SNAPIN, possibly via its association with CK2 and PKD kinases. CSN-dependent phosphorylation of TP53 and JUN promotes and protects degradation by the Ubl system, respectively.

# **Immunogen information**

Immunogen: COP9 constitutive photomorphogenic homolog subunit 4(Arabidopsis)

Synonyms: CSN4

Observed MW: 40; 46 kDa UniprotID: Q9BT78

## **Application**

1

#### Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)

Tel:(0086)027-87384275 Fax: (0086)027-87800889 www.fn-test.com




Reactivity: Human, Mouse, Rat


Tested Application: ELISA, WB, IHC, IP

Recommended dilution: WB: 1:500-1:2000; IP: 1:200-1:1000; IHC: 1:20-1:200

Image:




Immunohistochemistry of paraffin-embedded human pancreas tissue slide using FNab01872(COPS4 Antibody) at dilution of 1:50



IP Result of anti-COPS4 (IP:FNab01872, 4ug; Detection:FNab01872 1:300) with PC-3 cells lysate 1800ug.

# Wuhan Fine Biotech Co., Ltd.





A2780 cells were subjected to SDS PAGE followed by western blot with FNab01872(COPS4 antibody) at dilution of 1:500