

anti- CYP4F3 antibody

Product Information

Catalog No.:	FNab02173
Size:	100µg
Form:	liquid
Purification:	Immunogen affinity purified
Purity:	≥95% as determined by SDS-PAGE
Host:	Rabbit
Clonality:	polyclonal
Clone ID:	None
IsoType:	IgG
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months (Avoid repeated freeze / thaw cycles.)

Background

Isoform CYP4F3A: Catalyzes the omega-hydroxylation of leukotriene-B(4), a potent chemoattractant for polymorphonuclear leukocytes, it has low activity for arachidonic acid. Isoform CYP4F3B: Shows arachidonic acid omega-hydroxylase activity by mediating conversion of arachidonic acid to 20-hydroxyeicosatetraenoic acid(20-HETE)(PubMed:11461919, PubMed:16820285). Has a 30-fold higher Km for leukotriene-B(4) compared with CYP4F3A(PubMed:11461919). Also converts eicosapentaenoate(EPA) to 20-hydroxyeicosapentaenoate(20-OH-EPA)(PubMed:16820285). Acts as a polyunsaturated omega-3 fatty acids hydroxylase: mediates conversion of docosahexaenoate(DHA) to 22-hydroxydocosahexaenoate(PubMed:16820285).

Immunogen information

Immunogen:	cytochrome P450, family 4, subfamily F, polypeptide 3
Synonyms:	LTB4H
Observed MW:	60 kDa
UniprotID :	Q08477

Application

1


Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)

Tel :(0086)027-87384275

Fax: (0086)027-87800889 www.fn-test.com

Reactivity: Human
Tested Application: ELISA, WB
Recommended dilution: WB: 1:500-1:2000
Image:



HL-60 cells were subjected to SDS PAGE followed by western blot with FNab02173(CYP4F3 Antibody) at dilution of 1:300

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve. East Lake High-Tech Development Zone. Wuhan, Hubei, China(430206)

Tel : (0086)027-87384275

Fax: (0086)027-87800889 www.fn-test.com