

anti- EIF3D antibody

Product Information

Catalog No.: FNab02705

Size: 100μg Form: liquid

Purification: Protein A+G purification

Purity: ≥95% as determined by SDS-PAGE

Host: Mouse

Clonality: monoclonal

Clone ID: 2D11 IsoType: IgG1

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

Background

Component of the eukaryotic translation initiation factor 3(eIF-3) complex, which is required for several steps in the initiation of protein synthesis. The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl-tRNAi and eIF-5 to form the 43S pre-initiation complex(43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of post-termination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation. In case of FCV infection, plays a role in the ribosomal termination-reinitiation event leading to the translation of VP2(PubMed:18056426).

Immunogen information

Immunogen: eukaryotic translation initiation factor 3, subunit D

Synonyms: EIF3S7
Observed MW: 66 kDa
Uniprot ID: O15371

Application

1

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 Gaoxin Ave. East Lake High-Tech Development Zone. Wuhan, Hubei, China (430206)

Tel :(0086)027-87384275 Fax: (0086)027-87800889 <u>www.fn-test.com</u>




Reactivity: Human, Mouse, Rat


Tested Application: ELISA, WB, IHC

Recommended dilution: WB: 1:500-1:2000; IHC: 1:20-1:200

Image:



Immunohistochemistry of paraffin-embedded human brain using FNab02705(EIF3D antibody) at dilution of 1:50



HepG2 cells were subjected to SDS PAGE followed by western blot with FNab02705(EIF3D antibody) at dilution of 1:1000

Wuhan Fine Biotech Co., Ltd.