

anti- EMB antibody

Product Information

Catalog No.:	FNab02751
Size:	100µg
Form:	liquid
Purification:	Immunogen affinity purified
Purity:	\geq 95% as determined by SDS-PAGE
Host:	Rabbit
Clonality:	polyclonal
Clone ID:	None
IsoType:	IgG
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months (Avoid repeated freeze / thaw cycles.)

Background

Plays a role in the outgrowth of motoneurons and in the formation of neuromuscular junctions. Following muscle denervation, promotes nerve terminal sprouting and the formation of additional acetylcholine receptor clusters at synaptic sites without affecting terminal Schwann cell number or morphology. Delays the retraction of terminal sprouts following re-innervation of denervated endplates. May play a role in targeting the monocarboxylate transporters SLC16A1 and SLC16A7 to the cell membrane(By similarity).

Immunogen information

Immunogen:	embiginhomolog(Mouse)
Synonyms:	None
Observed MW:	70 kDa
UniprotID :	Q6PCB8

Application

Reactivity:	Human, Mouse
Tested Application:	ELISA, IHC, WB

Wuhan Fine Biotech Co., Ltd.


B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)

Tel :(0086)027-87384275 Fax: (0086)027-87800889 www.fn-test.com

1



Recommended dilution: WB: 1:500-1:2000; IHC: 1:20-1:200 Image:



Immunohistochemistry of paraffin-embedded human testis tissue slide using FNab02751(EMB Antibody) at dilution of 1:50

HeLa cells were subjected to SDS PAGE followed by western blot with FNab02751(EMB Antibody) at dilution of 1:1000

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)

Tel :(0086)027-87384275 Fax: (0086)027-87800889 <u>www.fn-test.com</u>