

anti- Endostatin antibody

Product Information

Catalog No.: FNab02762

Size: 100μg Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

Clone ID: None IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

Background

Endostatin is a naturally-occurring 20-kDa C-terminal fragment derived from COL18A1(also named as KNO). It is reported to serve as an anti-angiogenic agent, similar to angiostatin and thrombospondin. It potently inhibits endothelial cell proliferation and inhibits endothelial cell angiogenesis by binding to the heparan sulfate proteoglycans involved in growth factor signaling. Endostatin is a broad spectrum angiogenesis inhibitor and may interfere with the pro-angiogenic action of growth factors such as basic fibroblast growth factor(bFGF/FGF-2) and vascular endothelial growth factor(VEGF).

Immunogen information

Immunogen: collagen, type XVIII, alpha 1

Synonyms: Endostatin

Observed MW: 19 kDa, 27 kDa

Uniprot ID: P39060

Application

Reactivity: Human

1

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 Gaoxin Ave. East Lake High-Tech Development Zone. Wuhan, Hubei, China (430206)


Tel:(0086)027-87384275 Fax: (0086)027-87800889 <u>www.fn-test.com</u>




Tested Application: ELISA, WB, IHC

Recommended dilution: WB: 1:200-1:1000; IHC: 1:20-1:200

Image:



Immunohistochemistry of paraffin-embedded human liver using FNab02762(Endostatin, COL18A1 antibody) at dilution of 1:50



human liver tissue were subjected to SDS PAGE followed by western blot with FNab02762(Endostatin, COL18A1 antibody) at dilution of 1:500