

# anti- Histone H2A.X antibody

#### **Product Information**

Catalog No.: FNab03885

Size: 100μg Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

Clone ID: None IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

## **Background**

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene encodes a replication-independent histone that is a member of the histone H2A family, and generates two transcripts through the use of the conserved stem-loop termination motif, and the polyA addition motif.

#### **Immunogen information**

Immunogen: H2A histone family, member X

Synonyms: H2A histone family, member X, H2A.X, H2A/X, H2AFX, H2AX,

Histone H2A.x

Observed MW: 16 kDa UniprotID: P16104

## **Application**

1

### Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)


Tel :( 0086)027-87384275 Fax: (0086)027-87800889 <u>www.fn-test.com</u>




Reactivity: Human, Mouse, Rat Tested Application: ELISA, WB, IHC, IF

Recommended dilution: WB: 1:500 - 1:2000; IHC: 1:100 - 1:200; IF: 1:50 - 1:200

Image:



Immunohistochemistry of paraffin-embedded human breast cancer using FNab03885(H2AFX antibody) at dilution of 1:100



HL-60 cells were subjected to SDS PAGE followed by western blot with FNab03885(H2AFX antibody) at dilution of 1:1000