

anti- KDM2A antibody

Product Information

Catalog No.:	FNab04509
Size:	100µg
Form:	liquid
Purification:	Immunogen affinity purified
Purity:	≥95% as determined by SDS-PAGE
Host:	Rabbit
Clonality:	polyclonal
IsoType:	IgG
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months (Avoid repeated freeze / thaw cycles.)

Background

Histone demethylase that specifically demethylates 'Lys-36' of histone H3, thereby playing a central role in histone code. Preferentially demethylates dimethylated H3 'Lys-36' residue while it has weak or no activity for mono-and tri-methylated H3 'Lys-36'. May also recognize and bind to some phosphorylated proteins and promote their ubiquitination and degradation. Required to maintain the heterochromatic state. Associates with centromeres and represses transcription of small non-coding RNAs that are encoded by the clusters of satellite repeats at the centromere. Required to sustain centromeric integrity and genomic stability, particularly during mitosis.

Immunogen information

Immunogen:	lysine(K)-specific demethylase 2A
Synonyms:	CXXC8, FBL7, FBXL11, JHDM1A, KIAA1004
Observed MW:	80-132 kDa
Uniprot ID :	Q9Y2K7

Application

Reactivity:	Human, Mouse, Rat
Tested Application:	ELISA, WB, IHC
Recommended dilution:	WB: 1:500 - 1:2000; IHC: 1:50 - 1:200
Image:	

1


Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 Gaoxin Ave. East Lake High-Tech Development Zone. Wuhan, Hubei, China(430206)


Tel : (0086)027-87384275

Fax: (0086)027-87800889


www.fn-test.com



Immunohistochemistry of paraffin-embedded rat brain tissue slide using FNab04509(KDM2A Antibody) at dilution of 1:100



Jurkat cells were subjected to SDS PAGE followed by western blot with FNab04509(KDM2A Antibody) at dilution of 1:1000



NIH/3T3 cells were subjected to SDS PAGE followed by western blot with FNab04509(KDM2A Antibody) at dilution of 1:1000