

anti- MAGT1 antibody

Product Information

| | |
|---------------|---|
| Catalog No.: | FNab04955 |
| Size: | 100µg |
| Form: | liquid |
| Purification: | Immunogen affinity purified |
| Purity: | ≥95% as determined by SDS-PAGE |
| Host: | Rabbit |
| Clonality: | polyclonal |
| Clone ID: | None |
| IsoType: | IgG |
| Storage: | PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months (Avoid repeated freeze / thaw cycles.) |

Background

May be involved in N-glycosylation through its association with N-oligosaccharyltransferase. May be involved in Mg(2+) transport in epithelial cells.

Immunogen information

| | |
|--------------|-------------------------|
| Immunogen: | magnesium transporter 1 |
| Synonyms: | IAG2 |
| Observed MW: | 45-47 kDa |
| UniprotID : | Q9H0U3 |

Application


| | |
|-----------------------|---|
| Reactivity: | Human, Mouse, Rat |
| Tested Application: | ELISA, WB, IHC, IF |
| Recommended dilution: | WB: 1:500-1:2000; IHC: 1:20-1:200; IF: 1:20-1:200 |
| Image: | |

Wuhan Fine Biotech Co., Ltd.


B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve. East Lake High-Tech Development Zone. Wuhan, Hubei, China(430206)

Tel : (0086)027-87384275

Fax: (0086)027-87800889 www.fn-test.com



Immunohistochemistry of paraffin-embedded human liver using FNab04955(MAGT1 antibody) at dilution of 1:50



Jurkat cells were subjected to SDS PAGE followed by western blot with FNab04955(MAGT1 antibody) at dilution of 1:1000

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve. East Lake High-Tech Development Zone. Wuhan, Hubei, China(430206)

Tel : (0086)027-87384275

Fax: (0086)027-87800889 www.fn-test.com