

anti- MED30 antibody

Product Information

Catalog No.:	FNab05095
Size:	100µg
Form:	liquid
Purification:	Immunogen affinity purified
Purity:	≥95% as determined by SDS-PAGE
Host:	Rabbit
Clonality:	polyclonal
Clone ID:	None
IsoType:	IgG
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months (Avoid repeated freeze / thaw cycles.)

Background

Component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors.

Immunogen information

Immunogen:	mediator complex subunit 30
Synonyms:	THRAP6, TRAP25
Observed MW:	20-25 kDa
UniprotID :	Q96HR3

Application

Reactivity:	Human, Mouse
Tested Application:	ELISA, WB, IP

Wuhan Fine Biotech Co., Ltd.


B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve. East Lake High-Tech Development Zone. Wuhan, Hubei, China(430206)

Tel : (0086)027-87384275


Fax: (0086)027-87800889 www.fn-test.com

Recommended dilution: WB: 1:500-1:2000; IP: 1:200-1:2000

Image:



IP Result of anti-MED30 (IP:FNab05095, 4ug;
Detection:FNab05095 1:500) with mouse heart
tissue lysate 4000ug.



human heart tissue were subjected to SDS PAGE
followed by western blot with FNab05095(MED30
antibody) at dilution of 1:200