

anti- MEK1 antibody

Product Information

Catalog No.: FNab05112

Size: 100μg Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

Clone ID: None IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

Background

MEK1 is also named as MAP2K1(mitogen-activated protein kinase kinase 1), MAPKK1, PRKMK1, MKK1 and belongs to the MAP kinase kinase subfamily. It is a dual-specificity kinase that mediate ERK1 and ERK2 activation during adhesion and growth factor signaling(PMID:19219045). It also plays an essential role in extra-embryonic ectoderm during placentogenesis. MEK1 has 2 isoforms produced by alternative splicing with the molecular weight of 43 kDa and 41 kDa. It can be phosphorylated.

Immunogen information

Immunogen: mitogen-activated protein kinase kinase 1

Synonyms: ERK activator kinase 1, MAP kinase kinase 1, MAP2K1, MAPK/ERK

kinase 1, MAPKK 1, MAPKK1, MEK 1, MEK1, MKK1, PRKMK1

Observed MW: 40 kDa
Uniprot ID: Q02750

Application

Reactivity: Human, Mouse, Rat

1

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 Gaoxin Ave. East Lake High-Tech Development Zone. Wuhan, Hubei, China (430206)


Tel :(0086)027-87384275 Fax: (0086)027-87800889 <u>www.fn-test.com</u>




Tested Application: ELISA, WB, IHC, IF

Recommended dilution: WB: 1:500-1:2000; IHC: 1:50-1:200

Image:



Immunohistochemistry of paraffin-embedded rat lung using FNab05112(MEK1 antibody) at dilution of 1:100



mouse brain were subjected to SDS PAGE followed by western blot with FNab05112(MEK1 antibody) at dilution of 1:1000

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 Gaoxin Ave. East Lake High-Tech Development Zone. Wuhan, Hubei, China (430206)

Tel:(0086)027-87384275 Fax: (0086)027-87800889 <u>www.fn-test.com</u>