

anti- MICAL2 antibody

Product Information

Catalog No.:	FNab05178
Size:	100µg
Form:	liquid
Purification:	Immunogen affinity purified
Purity:	≥95% as determined by SDS-PAGE
Host:	Rabbit
Clonality:	polyclonal
Clone ID:	None
IsoType:	IgG
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months (Avoid repeated freeze / thaw cycles.)

Background

Nuclear monoxygenase that promotes depolymerization of F-actin by mediating oxidation of specific methionine residues on actin and regulates the SRF signaling. Acts by modifying nuclear actin subunits through the addition of oxygen to form methionine-sulfoxide, leading to promote actin filament severing and prevent repolymerization. Acts as a key regulator of the SRF signaling pathway elicited by nerve growth factor and serum: mediates oxidation and subsequent depolymerization of nuclear actin, leading to increase MKL1/MRTF-A presence in the nucleus and promote SRF:MKL1/MRTF-A-dependent gene transcription. Does not activate SRF:MKL1/MRTF-A through RhoA.

Immunogen information

Immunogen:	microtubule associated monoxygenase, calponin and LIM domain containing 2
Synonyms:	KIAA0750, MICAL2PV1, MICAL2PV2
Observed MW:	95, 112 kDa
UniprotID :	O94851

Application

1


Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)

Tel : (0086)027-87384275

Fax: (0086)027-87800889 www.fn-test.com

Reactivity: Human, Mouse, Rat
Tested Application: ELISA, IHC, IF
Recommended dilution: IHC: 1:20-1:200; IF: 1:20-1:200
Image:



Immunohistochemistry of paraffin-embedded human stomach cancer slide using FNab05178(MICAL2 Antibody) at dilution of 1:50