

anti- NR1H3 antibody

Product Information

Catalog No.: FNab05836

Size: 100μg Form: liquid

Purification: Protein A+G purification

Purity: ≥95% as determined by SDS-PAGE

Host: Mouse

Clonality: monoclonal

Clone ID: 7D2 IsoType: IgG2a

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

Background

Nuclear receptor. Interaction with RXR shifts RXR from its role as a silent DNA-binding partner to an active ligand-binding subunit in mediating retinoid responses through target genes defined by LXRES. LXRES are DR4-type response elements characterized by direct repeats of two similar hexanuclotide half-sites spaced by four nucleotides. Plays an important role in the regulation of cholesterol homeostasis, regulating cholesterol uptake through MYLIP-dependent ubiquitination of LDLR, VLDLR and LRP8. Interplays functionally with RORA for the regulation of genes involved in liver metabolism(By similarity). Exhibits a ligand-dependent transcriptional activation activity(PubMed:25661920).

Immunogen information

Immunogen: nuclear receptor subfamily 1, group H, member 3

Synonyms: Liver X receptor alpha, LXR a, LXRA, NR1H3, Oxysterols receptor

LXR alpha, RLD 1

Observed MW: 48 kDa Uniprot ID: Q13133

Application

1

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 Gaoxin Ave. East Lake High-Tech Development Zone. Wuhan, Hubei, China (430206)

Tel :(0086)027-87384275 Fax: (0086)027-87800889 <u>www.fn-test.com</u>




Reactivity: Human, Rat


Tested Application: ELISA, WB, IF

Recommended dilution: WB: 1:200-1:1000; IF: 1:10-1:100

Image:



Immunofluorescent analysis of HepG2 cells using FNab05836(NR1H3 Antibody) at dilution of 1:25 and Alexa Fluor 488-congugated AffiniPure Goat Anti-Mouse IgG(H+L)



MCF7 cells were subjected to SDS PAGE followed by western blot with FNab05836(NR1H3 antibody) at dilution of 1:500