

anti- PKM2-specific antibody

Product Information

Catalog No.:	FNab06495
Size:	100µg
Form:	liquid
Purification:	Immunogen affinity purified
Purity:	≥95% as determined by SDS-PAGE
Host:	Rabbit
Clonality:	polyclonal
Clone ID:	None
IsoType:	IgG
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months (Avoid repeated freeze / thaw cycles.)

Background

PKM2, also named as OIP3, PK2, PK3, PKM, p58, THBP1, CTHBP and Tumor M2-PK, belongs to the pyruvate kinase family. It is glycolytic enzyme that catalyzes the transfer of a phosphoryl group from phosphoenolpyruvate (PEP) to ADP, generating ATP. It stimulates POU5F1-mediated transcriptional activation. PKM2 plays a general role in caspase independent cell death of tumor cells. The activity of the M2 isoform (but not the M1 isoform) can be inhibited by tyrosine kinase signalling in tumour cells. The primary pyruvate kinase isoform before tumour development is PK-M1; however, the primary isoform from four independent tumours is PK-M2. (PMID:18337823) PKM2, Pyruvate kinase isozymes M1/M2, has 2 isoforms. The immunogen of this antibody is M2 isoform, also called PKM2. This antibody is specific to PKM2 isoform.

Immunogen information

Immunogen:	PKM2 pyruvate kinase, muscle
Synonyms:	CTHBP, OIP 3, OIP3, Opa interacting protein 3, p58, PK2, PK3, PKM, PKM2, PK-M2, PKM2 pyruvate kinase, muscle, Pyruvate kinase 2/3, Pyruvate kinase isozymes M1/M2, Pyruvate kinase muscle isozyme, TCB, THBP1, Tumor M2 PK
Observed MW:	58 kDa

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 Gaoxin Ave. East Lake High-Tech Development Zone. Wuhan, Hubei, China (430206)

Tel : (0086)027-87384275

Fax: (0086)027-87800889 www.fn-test.com

UniprotID : P14618


Application

Reactivity: Human, Mouse


Tested Application: ELISA, WB, IHC, IF, IP

Recommended dilution: WB: 1:1000-1:4000; IP: 1:200-1:1000; IHC: 1:100-1:400; IF: 1:10-1:100

Image:



Immunohistochemistry of paraffin-embedded human lung cancer tissue slide using FNab06495(PKM2-specific Antibody) at dilution of 1:200




Immunofluorescent analysis of MCF-7 cells, using FNab06495(PKM2-specific Antibody) at 1:25 dilution and Rhodamine-labeled goat anti-rabbit IgG (red)

Wuhan Fine Biotech Co., Ltd.


B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve. East Lake High-Tech Development Zone. Wuhan, Hubei, China(430206)

Tel : (0086)027-87384275

Fax: (0086)027-87800889 www.fn-test.com



IP Result of anti-PKM2-specific (IP:FNab06495, 4ug; Detection:FNab06495 1:300) with MCF-7 cells lysate 3250ug.



MCF7 cells were subjected to SDS PAGE followed by western blot with FNab06495(PKM2-specific antibody) at dilution of 1:1500