

# anti- PPARA antibody

#### **Product Information**

Catalog No.: FNab06663

Size: 100μg Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

# **Background**

Ligand-activated transcription factor. Key regulator of lipid metabolism. Activated by the endogenous ligand 1-palmitoyl-2-oleoyl-sn-glycerol-3-phosphocholine(16:0/18:1-GPC). Activated by oleylethanolamide, a naturally occurring lipid that regulates satiety. Receptor for peroxisome proliferators such as hypolipidemic drugs and fatty acids. Regulates the peroxisomal beta-oxidation pathway of fatty acids. Functions as transcription activator for the ACOX1 and P450 genes. Transactivation activity requires heterodimerization with RXRA and is antagonized by NR2C2. May be required for the propagation of clock information to metabolic pathways regulated by PER2.

# Immunogen information

Immunogen: peroxisome proliferator-activated receptor alpha

Synonyms: hPPAR, NR1C1, PPAR, PPAR alpha, PPARA

Observed MW: 52-55 kDa Uniprot ID: Q07869

### **Application**

Reactivity: Human, Mouse, Rat

#### 1

#### Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 Gaoxin Ave. East Lake High-Tech Development Zone. Wuhan, Hubei, China(430206)

Tel :( 0086)027-87384275 Fax: (0086)027-87800889 <u>www.fn-test.com</u>




Tested Application: ELISA, WB, IHC,IP

Recommended dilution: WB: 1:200-1:1000; IP: 1:200-1:1000; IHC: 1:50-1:200

Image:



Immunohistochemistry of paraffin-embedded human skeletal muscle tissue slide using FNab06663(PPARA Antibody) at dilution of 1:100



U-937 cells were subjected to SDS PAGE followed by western blot with FNab06663(PPARA antibody) at dilution of 1:300