

anti- PPAR δ antibody

Product Information

Catalog No.:	FNab06665
Size:	100 μ g
Form:	liquid
Purification:	Protein A+G purification
Purity:	\geq 95% as determined by SDS-PAGE
Host:	Mouse
Clonality:	monoclonal
Clone ID:	9B6
IsoType:	IgG1
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months (Avoid repeated freeze / thaw cycles.)

Background

Ligand-activated transcription factor. Receptor that binds peroxisome proliferators such as hypolipidemic drugs and fatty acids. Has a preference for poly-unsaturated fatty acids, such as gamma-linoleic acid and eicosapentanoic acid. Once activated by a ligand, the receptor binds to promoter elements of target genes. Regulates the peroxisomal beta-oxidation pathway of fatty acids. Functions as transcription activator for the acyl-CoA oxidase gene. Decreases expression of NPC1L1 once activated by a ligand.

Immunogen information

Immunogen:	peroxisome proliferator-activated receptor delta
Synonyms:	NPPARD, PPAR beta, NUCII, FAAR, PPAR delta
Observed MW:	54 kDa
Uniprot ID :	Q03181

Application

Reactivity:	Human, Mouse
Tested Application:	ELISA, WB

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 Gaoxin Ave. East Lake High-Tech Development Zone. Wuhan, Hubei, China(430206)


Tel : (0086)027-87384275

Fax: (0086)027-87800889

www.fn-test.com

Recommended dilution: WB: 1:500-1:1000

Image:



COLO 320 cells were subjected to SDS PAGE followed by western blot with FNab06665(PPARD antibody) at dilution of 1:1000

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 Gaoxin Ave. East Lake High-Tech Development Zone. Wuhan, Hubei, China(430206)

Tel : (0086)027-87384275

Fax: (0086)027-87800889

www.fn-test.com