

anti- RBPJ antibody

Product Information

Catalog No.: FNab07196

Size: 100μg Form: liquid

Purification: Protein A+G purification

Purity: ≥95% as determined by SDS-PAGE

Host: Mouse

Clonality: monoclonal

Clone ID: 2A10 IsoType: IgG1

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

Background

Transcriptional regulator that plays a central role in Notch signaling, a signaling pathway involved in cell-cell communication that regulates a broad spectrum of cell-fate determinations. Acts as a transcriptional repressor when it is not associated with Notch proteins. When associated with some NICD product of Notch proteins(Notch intracellular domain), it acts as a transcriptional activator that activates transcription of Notch target genes. Probably represses or activates transcription via the recruitment of chromatin remodeling complexes containing histone deacetylase or histone acetylase proteins, respectively. Specifically binds to the immunoglobulin kappa-type J segment recombination signal sequence. Binds specifically to methylated DNA. Binds to the oxygen responsive element of COX4I2 and activates its transcription under hypoxia conditions(4% oxygen)(PubMed:23303788).

Immunogen information

Immunogen: recombination signal binding protein for immunoglobulin kappa J region

Synonyms: IGKJRB, IGKJRB1, RBPJK, RBPSUH

Observed MW: 60 kDa Uniprot ID: Q06330

1

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 Gaoxin Ave. East Lake High-Tech Development Zone. Wuhan, Hubei, China (430206)

Tel:(0086)027-87384275 Fax: (0086)027-87800889 <u>www.fn-test.com</u>




Application

Reactivity: Human

Tested Application: ELISA, WB

Recommended dilution: WB: 1:500-1:2000

Image:



HepG2 cells were subjected to SDS PAGE followed by western blot with FNab07196(RBPJ antibody) at dilution of 1:1000