

anti- ROCK2 antibody

Product Information

Catalog No.: FNab07378

Size: 100μg Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

Clone ID: None IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

Background

Protein kinase which is a key regulator of actin cytoskeleton and cell polarity. Involved in regulation of smooth muscle contraction, actin cytoskeleton organization, stress fiber and focal adhesion formation, neurite retraction, cell adhesion and motility via phosphorylation of ADD1, BRCA2, CNN1, EZR, DPYSL2, EP300, MSN, MYL9/MLC2, NPM1, RDX, PPP1R12A and VIM. Phosphorylates SORL1 and IRF4. Acts as a negative regulator of VEGF-induced angiogenic endothelial cell activation. Positively regulates the activation of p42/MAPK1-p44/MAPK3 and of p90RSK/RPS6KA1 during myogenic differentiation. Plays an important role in the timely initiation of centrosome duplication. Inhibits keratinocyte terminal differentiation. May regulate closure of the eyelids and ventral body wall through organization of actomyosin bundles. Plays a critical role in the regulation of spine and synaptic properties in the hippocampus. Plays an important role in generating the circadian rhythm of the aortic myofilament Ca(2+) sensitivity and vascular contractility by modulating the myosin light chain phosphorylation.

Immunogen information

Immunogen: Rho-associated, coiled-coil containing protein kinase 2

Synonyms: KIAA0619
Observed MW: 161 kDa
UniprotID: 075116

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)

Tel:(0086)027-87384275 Fax: (0086)027-87800889 www.fn-test.com

1



Application


Reactivity: Human, Mouse, Rat

Tested Application: ELISA, WB, IHC, IF, IP


Recommended dilution: WB: 1:1000-1:10000; IP: 1:200-1:2000;; IHC: 1:20-1:200; IF: 1:10-

1:100

Image:



Immunohistochemistry of paraffin-embedded human liver using FNab07378(ROCK2 antibody) at dilution of 1:100



70kd


IP Result of anti-ROCK2(middle) (IP:FNab07378, 5ug; Detection:FNab07378 1:700) with HeLa cells lysate 1320ug.

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve. East Lake High-Tech Development Zone. Wuhan, Hubei, China (430206)

Tel: (0086)027-87384275 Fax: (0086)027-87800889 www.fn-test.com





HT-1080 cells were subjected to SDS PAGE followed by western blot with FNab07378(ROCK2 antibody) at dilution of 1:1000

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve. East Lake High-Tech Development Zone. Wuhan, Hubei, China (430206)

Tel: (0086)027-87384275 Fax: (0086)027-87800889 www.fn-test.com