

anti- SCARA5 antibody

Product Information

Catalog No.: FNab07623

Size: 100μg Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

Clone ID: None IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

Background

Ferritin receptor that mediates non-transferrin-dependent delivery of iron. Mediates cellular uptake of ferritin-bound iron by stimulating ferritin endocytosis from the cell surface with consequent iron delivery within the cell. Delivery of iron to cells by ferritin is required for the development of specific cell types, suggesting the existence of cell type-specific mechanisms of iron traffic in organogenesis, which alternatively utilize transferrin or non-transferrin iron delivery pathways. Ferritin mediates iron uptake in capsule cells of the developing kidney. Binds preferrentially ferritin light chain(FTL) compared to heavy chain(FTH1)(By similarity).

Immunogen information

Immunogen: scavenger receptor class A, member 5(putative)

Synonyms: None
Observed MW: 43 kDa
Uniprot ID: Q6ZMJ2

Application

Reactivity: Human, Mouse

1

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 Gaoxin Ave. East Lake High-Tech Development Zone. Wuhan, Hubei, China (430206)


Tel:(0086)027-87384275 Fax: (0086)027-87800889 <u>www.fn-test.com</u>



Tested Application: ELISA, WB

Recommended dilution: WB: 1:500-1:2000

Image:



mouse liver tissue were subjected to SDS PAGE followed by western blot with FNab07623(SCARA5 antibody) at dilution of 1:300

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 Gaoxin Ave. East Lake High-Tech Development Zone. Wuhan, Hubei, China (430206)

Tel: (0086)027-87384275 Fax: (0086)027-87800889 www.fn-test.com