

anti- SMYD2 antibody

Product Information

Catalog No.: FNab08048

Size: 100μg Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

Clone ID: None IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

Background

Protein-lysine N-methyltransferase that methylates both histones and non-histone proteins, including p53/TP53 and RB1. Specifically methylates histone H3 'Lys-4'(H3K4me) and dimethylates histone H3 'Lys-36'(H3K36me2). Shows even higher methyltransferase activity on p53/TP53. Monomethylates 'Lys-370' of p53/TP53, leading to decreased DNA-binding activity and subsequent transcriptional regulation activity of p53/TP53. Monomethylates RB1 at 'Lys-860'.

Immunogen information

Immunogen: SET and MYND domain containing 2

Synonyms: KMT3C
Observed MW: 50 kDa
UniprotID: Q9NRG4

Application

Reactivity: Human, Mouse

Tested Application: ELISA, WB, IF, IHC

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)


Tel:(0086)027-87384275 Fax: (0086)027-87800889 www.fn-test.com

1




Recommended dilution: WB: 1:500-1:2000; IHC: 1:20-1:200; IF: 1:20-1:200

Image:



Immunohistochemistry of paraffin-embedded human heart using FNab08048(SMYD2 antibody) at dilution of 1:100



mouse skeletal muscle tissue were subjected to SDS PAGE followed by western blot with FNab08048(SMYD2 antibody) at dilution of 1:500