

anti- STAT2 antibody

Product Information

Catalog No.: FNab08296

Size: 100μg Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

Clone ID: None IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

Background

Signal transducer and activator of transcription that mediates signaling by type I IFNs(IFN-alpha and IFN-beta). Following type I IFN binding to cell surface receptors, Jakkinases(TYK2 and JAK1) are activated, leading to tyrosine phosphorylation of STAT1 and STAT2. The phosphorylated STATs dimerize, associate with IRF9/ISGF3G to form a complex termed ISGF3 transcription factor, that enters the nucleus. ISGF3 binds to the IFN stimulated response element(ISRE) to activate the transcription of interferon stimulated genes, which drive the cell in an antiviral state(PubMed:9020188, PubMed:23391734). Acts as a regulator of mitochondrial fission by modulating the phosphorylation of DNM1L at 'Ser-616' and 'Ser-637' which activate and inactivate the GTPase activity of DNM1L respectively(PubMed:26122121).

Immunogen information

Immunogen: signal transducer and activator of transcription 2, 113kDa

Synonyms: None
Observed MW: 113 kDa
UniprotID: P52630

Application

1

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)


Tel:(0086)027-87384275 Fax: (0086)027-87800889 www.fn-test.com




Reactivity: Human, Mouse, Rat
Tested Application: ELISA, WB, IHC, IP

Recommended dilution: WB: 1:500-1:2000; IP: 1:200-1:1000; IHC: 1:20-1:200

Image:




Immunohistochemistry of paraffin-embedded human breast cancer using FNab08296(STAT2 antibody) at dilution of 1:50



IP Result of anti-STAT2 (IP:FNab08296, 4ug; Detection:FNab08296 1:500) with HeLa cells lysate 2800ug.

Wuhan Fine Biotech Co., Ltd.





HeLa cells were subjected to SDS PAGE followed by western blot with FNab08296(STAT2 antibody) at dilution of 1:500