

anti- TAF9 antibody

Product Information

Catalog No.:	FNab08489
Size:	100µg
Form:	liquid
Purification:	Immunogen affinity purified
Purity:	\geq 95% as determined by SDS-PAGE
Host:	Rabbit
Clonality:	polyclonal
Clone ID:	None
IsoType:	IgG
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months (Avoid repeated freeze / thaw cycles.)

Background

Essential for cell viability. TAF9 and TAF9B are involved in transcriptional activation as well as repression of distinct but overlapping sets of genes. May have a role in gene regulation associated with apoptosis. TAFs are components of the transcription factor IID(TFIID) complex, the TBP-free TAFII complex(TFTC), the PCAF histone acetylase complex and the STAGA transcription coactivator-HAT complex. TFIID or TFTC are essential for the regulation of RNA polymerase II-mediated transcription.

Immunogen information

Immunogen:	TAF9 RNA polymerase II, TATA box binding protein(TBP)-associated factor, 32kDa
Synonyms:	TAF2G, TAFII31
Observed MW:	24-28 kDa
UniprotID :	Q16594

Application

Reactivity: Human, Mouse, Rat


Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)

Tel :(0086)027-87384275 Fax: (0086)027-87800889 www.fn-test.com



Tested Application: ELISA, WB, IP Recommended dilution: WB: 1:500-1:2000; IP: 1:200-1:2000 Image:



IP Result of anti-TAF9 (IP:FNab08489, 4ug; Detection:FNab08489 1:1000) with mouse skeletal muscle tissue lysate 2200ug.

HeLa cells were subjected to SDS PAGE followed by western blot with FNab08489(TAF9 Antibody) at dilution of 1:1000

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 GaoxinAve.East Lake High-Tech Development Zone.Wuhan, Hubei, China(430206)

Tel :(0086)027-87384275 Fax: (0086)027-87800889 www.fn-test.com