

anti- TTF2 antibody

Product Information

Catalog No.:	FNab09092
Size:	100µg
Form:	liquid
Purification:	Immunogen affinity purified
Purity:	≥95% as determined by SDS-PAGE
Host:	Rabbit
Clonality:	polyclonal
Clone ID:	None
IsoType:	IgG
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months (Avoid repeated freeze / thaw cycles.)

Background

DsDNA-dependent ATPase which acts as a transcription termination factor by coupling ATP hydrolysis with removal of RNA polymerase II from the DNA template. May contribute to mitotic transcription repression. May also be involved in pre-mRNA splicing.

Immunogen information

Immunogen:	transcription termination factor, RNA polymerase II
Synonyms:	F2, HuF2, Lodestar homolog, Transcription release factor 2, TTF2
Observed MW:	130 kDa
Uniprot ID :	Q9UNY4

Application

Reactivity:	Human, Mouse, Rat
Tested Application:	ELISA, WB, IHC, IF, IP
Recommended dilution:	WB: 1:500-1:5000; IP: 1:200-1:1000; IF: 1:20-1:200; IHC: 1:20-1:200
Image:	

1


Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 Gaoxin Ave. East Lake High-Tech Development Zone. Wuhan, Hubei, China(430206)


Tel : (0086)027-87384275

Fax: (0086)027-87800889


www.fn-test.com



Immunohistochemistry of paraffin-embedded human ovary tumor using FNab09092(TTF2 antibody) at dilution of 1:100



IP Result of anti-TTF2 (IP:FNab09092, 4ug; Detection:FNab09092 1:500) with A2780 cells lysate 840ug.



A2780 cells were subjected to SDS PAGE followed by western blot with FNab09092(TTF2 antibody) at dilution of 1:500