

anti- TXNDC17 antibody

Product Information

Catalog No.:	FNab09124
Size:	100µg
Form:	liquid
Purification:	Immunogen affinity purified
Purity:	≥95% as determined by SDS-PAGE
Host:	Rabbit
Clonality:	polyclonal
Clone ID:	None
IsoType:	IgG
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months (Avoid repeated freeze / thaw cycles.)

Background

Disulfide reductase. May participate in various redox reactions through the reversible oxidation of its active center dithiol to a disulfide and catalyze dithiol-disulfide exchange reactions. Modulates TNF-alpha signaling and NF-kappa-B activation. Has peroxidase activity and may contribute to the elimination of cellular hydrogen peroxide.

Immunogen information

Immunogen:	thioredoxin domain containing 17
Synonyms:	TXNDC17, TXNL5, Thioredoxin like protein 5, TRP14, Protein 42 9 9
Observed MW:	14 kDa
Uniprot ID :	Q9BRA2

Application

Reactivity:	Human, Mouse, Rat
Tested Application:	ELISA, WB, IHC, IF
Recommended dilution:	WB: 1:500-1:2000; IF: 1:10-1:100; IHC: 1:20-1:200

Wuhan Fine Biotech Co., Ltd.


B9 Bld, High-Tech Medical Devices Park, No. 818 Gaoxin Ave. East Lake High-Tech Development Zone. Wuhan, Hubei, China(430206)

Tel : (0086)027-87384275


Fax: (0086)027-87800889

www.fn-test.com

Image:



Immunofluorescent analysis of HeLa cells, using TXNDC17 antibody FNab09124 at 1:25 dilution and Rhodamine-labeled goat anti-rabbit IgG (red).



Jurkat cells were subjected to SDS PAGE followed by western blot with FNab09124(TXNDC17 antibody) at dilution of 1:500