

anti- ZCCHC11 antibody

Product Information

Catalog No.:	FNab09609
Size:	100µg
Form:	liquid
Purification:	Immunogen affinity purified
Purity:	≥95% as determined by SDS-PAGE
Host:	Rabbit
Clonality:	polyclonal
Clone ID:	None
IsoType:	IgG
Storage:	PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 months (Avoid repeated freeze / thaw cycles.)

Background

Uridyltransferase that mediates the terminal uridylation of mRNAs with short (less than 25 nucleotides) poly(A) tails, hence facilitating global mRNA decay (PubMed:25480299). Involved in microRNA (miRNA)-induced gene silencing through uridylation of deadenylated miRNA targets. Also acts as a suppressor of miRNA biogenesis by mediating the terminal uridylation of some miRNA precursors, including that of let-7 (pre-let-7), miR107, miR-143 and miR-200c. Uridylated miRNAs are not processed by Dicer and undergo degradation. Degradation of pre-let-7 contributes to the maintenance of embryonic stem (ES) cell pluripotency (By similarity). Does not bind RNA directly, but recruited to RNA targets by RNA-binding protein LIN28A. Also catalyzes the 3' uridylation of miR-26A, a miRNA that targets IL6 transcript. This abrogates the silencing of IL6 transcript, hence promoting cytokine expression (By similarity). May also suppress Toll-like receptor-induced NF-kappa-B activation via binding to T2BP. Does not play a role in replication-dependent histone mRNA degradation. Due to functional redundancy between ZCCHC6 and ZCCHC11, the identification of the specific role of each of these proteins is difficult.

Immunogen information

Immunogen: zinc finger, CCHC domain containing 11

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 Gaoxin Ave. East Lake High-Tech Development Zone. Wuhan, Hubei, China (430206)

Tel : (0086)027-87384275

Fax: (0086)027-87800889

www.fn-test.com

Synonyms: KIAA0191, PAPD3, Terminal uridylyltransferase 4, TUT4, TUTase 4, ZCCHC11
Observed MW: 185 kDa
Uniprot ID : Q5TAX3


Application

Reactivity: Human, Mouse


Tested Application: ELISA, WB, IF, IP

Recommended dilution: WB: 1:200-1:2000; IP: 1:500-1:5000; IF: 1:20-1:200

Image:



IP Result of anti-ZCCHC11 (IP:FNab09609, 5ug; Detection:FNab09609 1:1000) with MCF-7 cells lysate 2000ug.



MCF7 cells were subjected to SDS PAGE followed by western blot with FNab09609(ZCCHC11 antibody) at dilution of 1:500