

# anti- PKC gamma antibody

#### **Product Information**

Catalog No.: FNab10007

Size:  $100\mu g$  Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

Clone ID: None IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

## **Background**

Protein kinase C(PKC) is a family of serine-and threonine-specific protein kinases that can be activated by calcium and second messenger diacylglycerol. PKC family members phosphorylate a wide variety of protein targets and are known to be involved in diverse cellular signaling pathways. PKC gamma is a neuron-specific member of the classical PKCs and is activated and translocated to subcellular regions as a result of various stimuli, including diacylglycerol synthesis, increased intracellular Ca(2+) and phorbol esters. Defects in this protein have been associated with spinocerebellar ataxia type 14(SCA14), an autosomal dominant neurodegenerative disease.

#### **Immunogen information**

Immunogen: protein kinase C, gamma

Synonyms: PKC gamma, PKCC, PKCG, PRKCG, Protein kinase C gamma type,

protein kinase C, gamma, SCA14

Observed MW: 78 kDa Uniprot ID: P63318

## **Application**

1

### Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 Gaoxin Ave. East Lake High-Tech Development Zone. Wuhan, Hubei, China (430206)

Tel:(0086)027-87384275 Fax: (0086)027-87800889 <u>www.fn-test.com</u>




Reactivity: Mouse, Rat


Tested Application: ELISA, WB, IHC

Recommended dilution: WB: 1:500-1:2000; IHC: 1:100-1:500

Image:



Immunohistochemistry of paraffin-embedded rat brain tissue slide using FNab10007(PRKCG Antibody) at dilution of 1:200



mouse brain tissue were subjected to SDS PAGE followed by western blot with FNab10007(PRKCG antibody) at dilution of 1:1000