

anti- MOK antibody

Product Information

Catalog No.: FNab10103

Size: 100μg Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

Background

This gene belongs to the MAP kinase superfamily. The gene was found to be regulated by caudal type transcription factor 2 (Cdx2) protein. The encoded protein, which is localized to epithelial cells in the intestinal crypt, may play a role in growth arrest and differentiation of cells of upper crypt and lower villus regions. Multiple alternatively spliced transcript variants encoding different isoforms have been observed for this gene.

Immunogen information

Immunogen: renal tumor antigen

Synonyms: RAGE, RAGE1

Observed MW: 40 kDa Uniprot ID: Q9UQ07

Application

Reactivity: Human, Rat

Tested Application: ELISA, WB, IHC

Recommended dilution: WB: 1:500-1:2000; IHC: 1:20-1:200

Image:


Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 Gaoxin Ave. East Lake High-Tech Development Zone. Wuhan, Hubei, China (430206)


Tel :(0086)027-87384275 Fax: (0086)027-87800889 <u>www.fn-test.com</u>

1





Immunohistochemistry of paraffin-embedded rat heart using FNab10103(MOK Antibody) at dilution of 1:100



HeLa cells were subjected to SDS PAGE followed by western blot with FNab10103(MOK antibody) at dilution of 1:1000