

anti- SUPT6H antibody

Product Information

Catalog No.: FNab10364

Size: 100μg Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

Clone ID: None IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

Background

Transcription elongation factor which binds histone H3 and plays a key role in the regulation of transcription elongation and mRNA processing. Enhances the transcription elongation by RNA polymerase II(RNAPII) and is also required for the efficient activation of transcriptional elongation by the HIV-1 nuclear transcriptional activator, Tat. Besides chaperoning histones in transcription, acts to transport and splice mRNA by forming a complex with IWS1 and the C-terminal domain(CTD) of the RNAPII subunit RPB1(POLR2A). The SUPT6H:IWS1:CTD complex recruits mRNA export factors(ALYREF/THOC4, EXOSC10) as well as histone modifying enzymes(such as SETD2), to ensure proper mRNA splicing, efficient mRNA export and elongation-coupled H3K36 methylation, a signature chromatin mark of active transcription. SUPT6H via its association with SETD1A, regulates both class-switch recombination and somatic hypermutation through formation of H3K4me3 epigenetic marks on activation-induced cytidine deaminase(AICDA) target loci. Promotes the activation of the myogenic gene program by entailing erasure of the repressive H3K27me3 epigenetic mark through stabilization of the chromatin interaction of the H3K27 demethylase KDM6A.

Immunogen information

Immunogen: suppressor of Ty 6 homolog(S. cerevisiae)

Synonyms: KIAA0162, SPT6H

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 Gaoxin Ave. East Lake High-Tech Development Zone. Wuhan, Hubei, China (430206)

Tel:(0086)027-87384275 Fax: (0086)027-87800889 <u>www.fn-test.com</u>

1



Observed MW: None

Uniprot ID: Q7KZ85


Application

Reactivity: Human

Tested Application: ELISA, IHC

Recommended dilution: IHC: 1:50 - 1:200

Image:



Immunohistochemistry of paraffin-embedded human kidney using FNab10364(SUPT6H antibody) at dilution of 1:100