

anti- SATB2 antibody

Product Information

Catalog No.: FNab10798

Size: 100μg Form: liquid

Purification: Immunogen affinity purified

Purity: ≥95% as determined by SDS-PAGE

Host: Rabbit

Clonality: polyclonal

IsoType: IgG

Storage: PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12

months (Avoid repeated freeze / thaw cycles.)

Background

Binds to DNA, at nuclear matrix-or scaffold-associated regions. Thought to recognize the sugar-phosphate structure of double-stranded DNA. Transcription factor controlling nuclear gene expression, by binding to matrix attachment regions(MARs) of DNA and inducing a local chromatin-loop remodeling. Acts as a docking site for several chromatin remodeling enzymes and also by recruiting corepressors(HDACs) or coactivators(HATs) directly to promoters and enhancers. Required for the initiation of the upper-layer neurons(UL1) specific genetic program and for the inactivation of deep-layer neurons(DL) and UL2 specific genes, probably by modulating BCL11B expression. Repressor of Ctip2 and regulatory determinant of corticocortical connections in the developing cerebral cortex. May play an important role in palate formation. Acts as a molecular node in a transcriptional network regulating skeletal development and osteoblast differentiation.

Immunogen information

Immunogen: SATB homeobox 2

Synonyms: DNA binding protein SATB2, KIAA1034, SATB homeobox 2, SATB2

Observed MW: 69-83 kDa Uniprot ID: Q9UPW6

1

Wuhan Fine Biotech Co., Ltd.

B9 Bld, High-Tech Medical Devices Park, No. 818 Gaoxin Ave. East Lake High-Tech Development Zone. Wuhan, Hubei, China (430206)

Tel :(0086)027-87384275 Fax: (0086)027-87800889 <u>www.fn-test.com</u>




Application

Reactivity: Human

Tested Application: ELISA, WB, IHC

Recommended dilution: WB: 1: 200-1: 2000; IHC: 1: 20-1: 200

Image:



MCF7 cells were subjected to SDS PAGE followed by western blot with FNab10798(SATB2 antibody) at dilution of 1:1000